

Hallux rigidus is a disorder of the joint located at the base of the big toe. It causes pain and stiffness in the joint, and with time it gets increasingly harder to bend the toe.

‘Hallux’ refers to the big toe, while ‘rigidus’ indicates that the toe is rigid and cannot move. Hallux rigidus is actually a form of degenerative arthritis.

This disorder can be very troubling and even disabling, since we use the big toe whenever we walk, stoop down, climb up, or even stand. Many patients confuse hallux rigidus with a bunion, which affects the same joint, but they are very different conditions requiring different treatment.

Because hallux rigidus is a progressive condition, the toe’s motion decreases as time goes on. In its earlier stage, when motion of the big toe is only somewhat limited, the condition is called “hallux limitus.” But as the problem advances, the toe’s range of motion gradually decreases until it potentially reaches the end stage of “rigidus,” in which the big toe becomes stiff, or what is sometimes called a “frozen joint.”

Causes

Common causes of hallux rigidus are faulty function (biomechanics) and structural abnormalities of the foot that can lead to osteoarthritis in the big toe joint. This type of arthritis – the kind that results from “wear and tear” – often develops in people who have defects that change the way their foot and big toe functions. For example, those with fallen arches or excessive pronation (rolling in) of the ankles are susceptible to developing hallux rigidus. In some people, hallux rigidus runs in the family and is a result of inheriting a foot type that is prone to developing this condition. In other cases, it is associated with overuse – especially among people engaged in activities or jobs that increase the stress on the big toe, such as workers who often have to stoop or squat. Hallux rigidus can also result from an injury, such as stubbing your toe. Or it may be caused by inflammatory diseases such as rheumatoid arthritis or gout. Your foot and ankle surgeon can determine the cause of your hallux rigidus and recommend the best treatment.


Symptoms

Early signs and symptoms include:
 Pain and stiffness in the big toe during use (walking, standing, bending, etc.)
 Pain and stiffness aggravated by cold, damp weather
 Difficulty with certain activities (running, squatting)
 Swelling and inflammation around the joint
 As the disorder gets more serious, additional symptoms may develop, including:
 Pain, even during rest
 Difficulty wearing shoes because bone spurs (overgrowths) develop

Dull pain in the hip, knee, or lower back due to changes in the way you walk
 Limping (in severe cases)

Diagnosis

The sooner this condition is diagnosed, the easier it is to treat. Therefore, the best time to see a foot and ankle surgeon is when you first notice symptoms. If you wait until bone spurs develop, your condition is likely to be more difficult to manage.



In diagnosing hallux rigidus, the surgeon will examine your feet and move the toe to determine its range of motion. X-rays help determine how much arthritis is present as well as to evaluate any bone spurs or other abnormalities that may have formed.

Non-Surgical Treatment

1. Shoes with a large toe box put less pressure on your toe. Stiff or rocker-bottom soles are recommended.
2. Orthotic devices. Custom orthotic devices may improve foot function.
3. Medications. Oral nonsteroidal anti-inflammatory drugs (NSAIDs), such as ibuprofen, may be recommended to reduce pain and inflammation.
4. Injection therapy. Injections of corticosteroids may reduce inflammation and pain.

When Is Surgery Needed?

When conservative options fail and deformity is affecting daily lifestyle, surgery may be indicated. Depending on your deformity, removal of the impeding bone spur to improve ROM is indicated in milder deformities. When the joint is at end stage, a partial implant or fusion of the great toe joint is indicated. The procedure of choice or combination of procedures is considered according to xray findings, your age, activity level, and other factors. The length of the recovery period will vary, depending on the procedure or procedures performed.

